

Spatial and temporal variation in macroparasite communities of three-spined stickleback

REBECCA E. YOUNG* and ANDREW D. C. MACCOLL

School of Life Sciences, University Park, University of Nottingham, Nottingham NG7 2RD, UK

(Received 18 May 2016; revised 13 September 2016; accepted 14 September 2016)

SUMMARY

Patterns in parasite community structure are often observed in natural systems and an important question in parasite ecology is whether such patterns are repeatable across time and space. Field studies commonly look at spatial or temporal repeatability of patterns, but they are rarely investigated in conjunction. We use a large dataset on the macroparasites of the three-spined stickleback, *Gasterosteus aculeatus* L., collected from 14 locations on North Uist, Scotland over an 8-year period to investigate: (1) repeatability of patterns in parasite communities among populations and whether variation is consistent across years, (2) whether variation between years can be explained by climatic variation and progression of the season and (3) whether variation in habitat characteristics explain population differences. Differences in relative abundance and prevalence across populations were observed in a number of parasites investigated indicating a lack of consistency across years in numerous parasite community measures; however, differences between populations in the prevalence and abundance of some parasites were consistent throughout the study. Average temperature did not affect parasite community, and progression of the season was only significant for two of 13 community measures. Two of the six habitat characteristics investigated (pH and calcium concentration) significantly affected parasite presence.

Key words: stickleback, parasite community, repeatability.

INTRODUCTION

A key goal of many scientific disciplines is the identification of general laws or principles based upon recurring and predictable patterns (Poulin, 2007). Such patterns can be used not only to formulate laws explaining observations in nature and their underlying mechanisms, but also as a basis for testable hypotheses (Lawton, 1999; Poulin, 2007). However, finding laws which can be applied in all cases is difficult in ecology as the complexity of natural systems results in identification of circumstantial patterns which are not applicable in all situations (Poulin, 2007). Many ecologists, including parasite ecologists, continue to search for repeatable patterns across time, geographical area and taxa (Poulin, 2007; Kennedy, 2009; de Roij and MacColl, 2012). There has been much uncertainty about the extent to which parasite communities are structured, as well as whether observed relationships are sustained or transient (Behnke *et al.* 2008a). Identifications of patterns in parasite occurrences may provide valuable insights into the shaping of parasite communities and interactions, as well as the dynamics of host–parasite relationships (Behnke, 2008; de Roij and MacColl, 2012).

The organization of parasite communities infecting a species is hierarchical and can be looked at on a number of levels, ranging from infracommunities,

through to component communities and finally the total parasite fauna (as defined by Bush *et al.* 1997). The different ecological processes acting at different levels influence how dynamic community structure is, with the lowest levels being most subject to temporal and spatial variation (Behnke *et al.* 2008a). At the component community level, numerous factors, both extrinsic (location, year and season) and intrinsic (host age, sex and resistance), can be important contributors to fluctuations commonly observed (Abu-Madi *et al.* 1998; Behnke *et al.* 2008a). Extrinsic factors contributing to community variation have been the focus of numerous studies looking at the effects of season (Bolek and Coggins, 2000), year (Kennedy *et al.* 2001) and population heterogeneities (Calvete *et al.* 2004). Despite a large body of work looking at temporal and spatial variation, it has been less common for these effects to be investigated in conjunction with each other (de Roij and MacColl, 2012). Patterns observed when looking at population or year/season alone provide a snapshot of community composition and, whilst they may succeed in uncovering patterns in community structure, these patterns are rarely consistent when spatially or temporally replicated communities are observed (González and Poulin, 2005). Thus, such patterns are likely to describe characteristics of a certain population at one time and place, rather than reflect the host's parasite community as a whole (Vidal-Martínez and Poulin, 2003). Kennedy (1997) emphasised the importance of long-term datasets in furthering our understanding

* Corresponding author: School of Life Sciences, University Park, University of Nottingham, Nottingham NG7 2RD, UK. E-mail: rebeccayoung1291@gmail.com

of parasite ecology; such datasets facilitate much needed investigation of repeatability of observed patterns across space and time.

Factors affecting parasite distribution may be viewed at two levels: the host and the environment in which the host resides (Cardon *et al.* 2011). Effects at the host level include intrinsic variables, such as age, body size, genetic susceptibility and sex (Behnke *et al.* 2001; Blanchet *et al.* 2009) although the relative significance of each of these factors is currently unclear (Wilson *et al.* 2002). To get a full understanding of parasite community dynamics, it is important to consider also biotic and abiotic factors correlated with observed variation, which can strongly affect community dynamics (Lively *et al.* 2014). These environmental contributors relate to the habitat in which hosts and parasites live: for example, host density, diet and climate (Cardon *et al.* 2011; de Roij and MacColl, 2012). These factors are suggested to play a role in shaping component communities either directly, by affecting free-living parasite stages, or indirectly, by affecting survival of intermediate hosts (Pietrock and Marcogliese, 2003). Previous spatial and temporal studies have incorporated abiotic factors into their work to determine whether they could explain variation in species richness, prevalence and abundance across study sites (Marcogliese and Cone, 1996; Goater *et al.* 2005; de Roij and MacColl, 2012). In this study, we use the spatio-temporal variation in parasite communities infecting three-spined sticklebacks, *Gasterosteus aculeatus* L. (hereafter referred to as stickleback), in 14 freshwater lochs on the Scottish island of North Uist to try to give insight into factors contributing to this variation. It continues from work started by de Roij and MacColl (2012), who found that parasite communities in 12 of these lochs remained constant over a 2-year period (2007 and 2008), but found that these patterns could not be explained by effects of limnological, physiochemical and geomorphological variation (pH, calcium concentration, chlorophyll A concentration, dissolved organic carbon and loch surface area) on occurrences of parasites.

There are numerous benefits to using the North Uist study system in assessing parasite spatio-temporal variation and environmental effects. Firstly, the island has a large network of lochs which, due to their geographic isolation, can be considered to contain separate populations of sticklebacks, typically with high population densities making it easy to collect sufficient sample sizes (de Roij and MacColl, 2012). Also, unlike many studies of spatial and temporal repeatability, this system is confined to a small spatial scale. This allows comparison of a number of different populations within a small geographic area, and thus a greater focus on the impact of local factors (de Roij and MacColl, 2012). Since the work of de Roij and

MacColl (2012), further data have been collected from these populations in 2011, 2013 and 2014, resulting in a large dataset which will be used to investigate: (i) parasite community composition and repeatability, (ii) possible explanations behind between-year variation, based on year-to-year temperature variation and seasonal impacts, and (iii) whether environmental variables can explain between-site variation. By considering these factors in models of parasite community measures we hope to be able to identify possible mechanisms explaining patterns of variation observed when looking at spatial and temporal variation.

Mechanistic explanations of variation

Climate has been found directly to affect the rate of parasite development and survival of transmission stages (Chappell, 1969; Behnke *et al.* 2005). Sampling point in the season can thus affect parasite occurrence, as observed by increased infection with diplostomid species in late spring (Pennycuik, 1971). Therefore the average temperature and the point in the season (Julian date) at which parasite data were collected were considered in analysis.

Six factors (geomorphological, biotic and abiotic), were included as correlates of spatial variation: loch surface area, mean depth, calcium concentration (Ca^{2+} conc.), pH, log *Pungitius pungitius* and stickleback catch rate. Previous work gives some indication that each of these factors may be of importance to parasite communities. Due to the expected species area relationship, loch surface area is of importance as larger water bodies would be expected to contain a higher parasite species richness (Connor and McCoy, 1979; Ebert *et al.* 2001). Mean loch depth is anticipated to be more important in determining measures of individual parasite prevalence, as habitat use by intermediate hosts affects where parasites may be found, e.g. diplostomids infect snails utilising the littoral zones and cestodes infect copepods in pelagic zones (Marcogliese and Cone, 1991). Calcium concentration, which is strongly positively correlated with pH (MacColl *et al.* 2013), has been found to affect the presence of diplostomids, perhaps because high calcium concentration is required to support snail intermediate hosts (Curtis and Rau, 1980). Similarly, in more acidic reservoirs, perch, *Perca fluviatilis*, have reduced species richness and an absence of all but one digenean species (Halmetoja *et al.* 2000). *Pungitius pungitius* (nine-spined stickleback), is a competitor of three-spined stickleback and a potential alternative host for a number of parasites, including *Protecephalus filicollis* and *Schistocephalus solidus* (Dartnall, 1973). *Pungitius pungitius* is found in 10 of the 14 lochs investigated in this study (see supplementary data Table S2); therefore, as host density can effect parasite transmission, *P. pungitius*

density (describing the density of nine-spined stickleback) and stickleback catch rate (a proxy for stickleback density) are also taken into consideration (Soleng *et al.* 1999; Arneberg, 2002).

MATERIALS AND METHODS

Fish populations, sampling and parasite identification

A total of 1130 stickleback were collected from 14 geographically isolated, freshwater lochs on North Uist, Scotland. Stickleback were sampled over a two week period during the breeding season (April–May) in 5 years between 2007 and 2014 (no relevant samples were collected in 2009, 2010 or 2012). Fish were collected using minnow traps (Gee traps, Dynamic Aqua, Vancouver). In general, 20–30 traps were set overnight and lifted the following day, spread out along the shoreline of the lochs and focused on areas with vegetation: where sticklebacks are more commonly found. Samples of at least 20 fish were selected haphazardly from those caught although in some instances the samples were smaller if 20 fish were not caught.

Fish were transferred from traps into polystyrene boxes, with an air stone, for transport and were stored in these boxes in lake water for a maximum of 48 h. Within this time (and normally within 24 h), fish were killed and thoroughly inspected for macroparasites under a dissection microscope. Parasites were identified (generally to species level, using a key for parasites of freshwater fish (Bykhovskaya-Pavloskaya *et al.* 1964)) and recorded, along with measurements of the standard length (to the nearest 0.1 mm) and weight of the whole fish (to the nearest 0.0001 g). First the caudal, dorsal and anal fins were inspected, then the rest of the body surface and the gills. The abundance of the ectoparasites present was recorded. In 2007 and 2008, only the left eye was removed and dissected: in all subsequent years, both eyes were dissected and lens and retinal tissue inspected for parasites. Data for the left eye was strongly correlated with that for both eyes combined for all three eye dwelling parasites (*Apatemon gracilis* $R = 0.940$, $P < 0.001$; *Diplostomum gasterostei* $R = 0.917$, $P < 0.001$; *Diplostomum spathaceum* $R = 0.983$, $P < 0.001$) so just left eye data are used in subsequent analysis. The body cavity was opened and any parasites present in the peritoneal cavity were identified and counted. Fish were labelled and preserved in 70% ethanol and dissection was completed after returning to the laboratory in Nottingham where intestines were removed and thoroughly checked. Where possible, parasites were identified to species level. Two cestodes found in the intestine, *Bothriocephalus scorpii* and *Eubothrium crassum*, were generally immature and are very difficult to differentiate at such an early stage in the life cycle (Andersen and Valtonen, 1990); thus, they were combined and recorded as a single ‘Cestoda gen. spp.’ count. It is

likely that in the present analysis of freshwater populations most of the cestodes in this grouping were *E. crassum*, since identifiable *B. scorpii* were only ever found in stickleback in saltwater (A. D. C. MacColl, personal observations).

Environmental data collection

Samples of fish were collected at slightly different times each year between late April and late May and year-to-year variation (probably in winter and spring weather) meant that the season had progressed to varying extents between years. Such variation could alter the proportion of fish in breeding condition, and the state of parasitic infections. To account for the extrinsic factor of climatic variation between years the average temperature during the months before each sample was collected were obtained via publically available Met Office UK climate data (<http://www.metoffice.gov.uk/climate/uk/stationdata/>). Using historic station data from Stornoway Airport (located on the Isle of Lewis, Scotland, approximately 82 km from North Uist), the average temperature for March and April was calculated for each year sampled. Variation in time of the year at which data were collected was accounted for using Julian date; indicating the time elapsed since January 1st.

Two abiotic factors, representing the dominant axis of water chemistry on North Uist (Waterston *et al.*, 1979) were measured for each loch. Measurements of pH were taken between April 2006 and May 2013 using a calibrated pH meter (Multi 340i, Semat International) and an average was taken from three to six readings. To measure calcium concentration, filtered water samples were collected in May 2011, and acidified with nitric acid before freezing and returning to the University of Nottingham for analysis using inductively coupled plasma mass spectrometry (ICP-MS, Thermo-Fisher XSeries^{II}). Mean stickleback catch rate was measured by ‘catch per unit effort’ (CPUE); the number of sticklebacks caught was divided by the number of traps set per night. The average of these measurements was then taken for two to 4 years between 2009 and 2013 to provide a mean stickleback catch rate. Density of the competitor species *P. pungitius* was calculated as the percentage of nine-spined stickleback, rather than three-spined stickleback, in a haphazard sample (minimum size = 100) of all stickleback caught. An average for these percentages was then taken from 3 years (2010, 2011 and 2013) and the natural log of these percentages was used for comparisons. Loch surface area was estimated using web-based planimeter software (<http://www.freemaptools.com/area-calculator.htm>) and Google Earth, and the mean depth was calculated from 30 readings of depth taken from a boat using a hand-held depth sounder (Platimo Echotest II) at various locations around lochs.

Methods for statistical analysis

In analyses of the patterns in parasite occurrence, a sample of 1130 fish was used (see supplementary data Table S1 for full details of samples). Data analysis was carried out using computer programmes GenStat (15th edition, VSN international Ltd, Hemel Hempstead, UK) and Microsoft Excel, 2010 (Microsoft Corporation, Washington, USA).

Parasite communities: general patterns. The following summary statistics were calculated for each population/year combination in order to establish general patterns of community composition: species richness, abundance and prevalence of parasites [as described by Bush *et al.* (1997)]. Prevalence and abundance are used in conjunction because, although not completely independent, nevertheless the two measures contain different information about the distribution of parasites across hosts, and allow contrasting inference about the likely effect of parasites on host populations (Anderson and May, 1987). As well as calculating prevalence for individual populations, presence/absence data were used to calculate the overall prevalence across all populations and years in order to quantify how commonly parasites occur and thus, determine which should be considered for further analysis. Parasites which failed to exceed an overall prevalence of 10% were not used in further analysis (MacColl, 2009). Simpson's diversity index (1-D) was also used as a simple measure of diversity at the component community level (Magurran, 2004).

Variation in abundance and individual prevalence of parasites: (i) the individual level. Univariate generalized linear models (GLMs) were used to analyse parasite abundance, individual prevalence and species richness at the level of the individual host. The 13 dependent variables modelled were species richness, and the prevalence and abundance of each of the six key parasite groups. Species richness was modelled using normal errors and an identity link function. Parasite prevalences were modelled using binomial errors and a logit link function: ('1' and '0' for infected and non-infected fish respectively). Parasite abundances were modelled using negative binomial errors and a logarithmic link function. Population, year and sex were included as explanatory variables for all models and standard length was fitted as a covariate. In the most complex model, a population \times year interaction term was included. The deletion approach was used to reach a minimum adequate model, whereby the most complex model was tested first and non-significant terms were sequentially removed. *P*-values were corrected throughout using a sequential Bonferroni correction to account for multiple comparisons. Results are displayed in tables including the estimates of coefficients for continuous data.

Variation in mean abundance and prevalence of parasites: (ii) the population level. Average species richness, parasite prevalences and mean parasite abundances were modelled as dependent variables across all years and all populations studied in order to find mechanistic explanations for any variation observed. The 13 dependent variables remained the same but distributions differed when average measurements were modelled. Prevalence was normally distributed, as average prevalences approximate to a normal distribution. Average abundances are not integers; therefore, it was no longer appropriate to use negative binomial distribution so average abundance was log transformed and a normal distribution used. In all cases, the average length and sex ratio were included as explanatory variables as body length is commonly observed in nature to correlate with parasite presence, especially in fish (Poulin, 1997) and sex can affect parasite infection (Behnke, 2008).

Temporal climatic and seasonal effects. Annual averages per population were calculated for parasite community measures, temperature and Julian date. GLMs were used to model annual averages of dependent variables for each population against average temperature and Julian date to look for the effects of climate and season, respectively.

Spatial environmental effects. To identify mechanistic explanations for variation between populations, GLMs were used to model overall population averages across all years of dependent variables against environmental variables. Mean pH, calcium concentration, stickleback catch rate and log (relative density *P. pungitius*) were used as explanatory variables in all models and loch surface area was included for community measures (species richness), whilst mean loch depth was used for parasite measures (abundance and prevalence). These two different measures were used because the area of water bodies has previously been shown to impact the number of parasite species present (Ebert *et al.* 2001) and different parasites can be found in different depths of water (Marcogliese and Cone, 1991).

RESULTS

Parasite communities

The component community of macroparasites infecting *G. aculeatus* consisted of 12 parasites (Table 1) with a total of 78% of fish being infected with at least one parasite ($n = 878$ out of 1130). Prevalence was calculated across all populations and years in order to identify commonly occurring parasites (Table 1). Seven parasite taxa were found to exceed 10% prevalence across samples: the crustacean *Thersitina gasterostei*, the monogenean

Table 1. Prevalence of all macroparasites of three-spined stickleback ($n = 1130$) identified in 14 populations on the Scottish island of North Uist across an 8-year period

Parasite	No. of fish infected	% Prevalence
Monogenea		
<i>Gyrodactylus arcuatus</i>	325	25.66 ^a
Digenea		
<i>Diplostomum gasterostei</i>	505	44.43 ^a
<i>Diplostomum spathaceum</i>	55	4.78
<i>Apatemon gracilis</i>	273	24.07 ^a
<i>Podocotyle</i> sp.	2	0.18
Arthropoda		
<i>Thersitina gasterostei</i>	136	12.04 ^a
Nematode		
Nematoda gen. sp.	7	0.62
Cestodes		
<i>Diphyllobothrium</i> sp.	98	8.67
<i>Schistocephalus solidus</i>	205	17.97 ^a
<i>Proteocephalus filicollis</i>	175	14.60 ^a
Cestoda gen. spp. ^b	124	10.97 ^a

^a Parasites exceeding average 10% prevalence across all populations and years sampled which were considered in further analysis.

^b This group consists of two species of cestodes found in the intestine *E. crassum* and *B. scorpii* which were generally too immature to differentiate. In these freshwater populations, the majority of these cestodes are likely to be *E. crassum*, since identifiable *B. scorpii* have only been found in stickleback living in saltwater on North Uist.

Gyrodactylus arcuatus, the trematodes *D. gasterostei* and *A. gracilis* and the cestodes *S. solidus* and *Proteocephalus filicollis*, and the group 'Cestoda gen. spp.', consisting of *B. scorpii* and *E. crassum*.

Most of these parasites are described as common parasites and considered for further analysis. *Thersitina gasterostei* only occurred in three populations and was found in fewer than 20% of 57 population samples collected from different lochs across 5 years, so was not included in further analysis.

Parasite communities: general patterns. Overall infection levels calculated across all years for each individual population were generally high: seven of the 14 populations had more than 80% of fish infected with at least one parasite (Fig. 1) and only one population had fewer than 60% of fish infected (Daim, 31.25%). Furthermore, infracommunities consisting of more than one parasite were found to be very common (Fig. 1): three populations had a large proportion (>80%) of fish infected with at least two parasites (Gill, 94%; Host, 83.5% and Reiv, 94.2%) and a further four had over 50% of fish infected with multiple parasites (Buai, 57.0%; Chru, 54.8%; Maga, 63.6% and Mora, 56.9%). Two populations (Gill, 84% and Reiv, 81.2%) had a large proportion of fish infected with at least three parasites. Mean species richness, calculated for each population, ranged from 0.45 ± 0.88 (Daim) to 3.7 ± 1.17 (Gill) (Fig. 2A). Parasite diversity (1-D) did not vary significantly between years and populations (Fig. 2B, Wald $F = 1.19$, D.F. = 4, 39, $P = 0.33$; Wald $F = 1.30$, D.F. = 13, 39, $P = 0.252$, respectively).

Variation in abundance and individual prevalence of parasites: (i) the individual host level. GLMs of the abundance and prevalence of parasites in individual hosts revealed some common patterns. Length had a significant effect on species richness, prevalence of all parasite species, apart from the three cestodes (*S. solidus*, *P. filicollis* and Cestoda gen. spp.) and abundance of all parasites apart from *G. arcuatus* and *S. solidus* (Table 2). Correlations were positive for all parasites, apart from *P. filicollis*, indicating a greater prevalence and higher abundance of parasites in larger fish (Table 2). Sex did not generally explain a significant proportion of the variance in either parasite abundance or prevalence, although it was significant in predicting the prevalence and abundance of *S. solidus*, both of which are greater in males than in females (Table 2). The sex ratio of samples collected ranged between 0.30 and 0.62 and all but two population samples were female biased. Buai had more males (sex ratio = 0.62) and Chru had equal numbers of males and females.

There was significant variation between populations for all response variables (Table 2). Parasite species richness, abundance and prevalence also varied between years, except for the prevalence and abundance of *P. filicollis* and the prevalence of *G. arcuatus* (Table 2). The year \times population interaction term was significant in a number of models: species richness, abundance (except for *P. filicollis* which was consistently very low in the majority of populations, see below [Fig. 3A]) and prevalence of *D. gasterostei* and *A. gracilis* (Table 2) were all significant, indicating that variation was not completely




Fig. 1. Percentage of three-spined stickleback in 14 North Uist lochs infected with at least one (white), two (grey) and three (black) parasites ($\pm 95\%$ confidence interval).




Fig. 2. Variation in diversity of parasites of three-spined stickleback in 14 different lochs on North Uist. (A) Average species richness per population (\pm S.E.); (B) Simpson's diversity index.

consistent within populations across years in these instances. This makes the interpretation of patterns of spatiotemporal variation difficult, but this can be clarified through the use of figures.

For example, the prevalence of *P. filicollis* was consistently below 20% in the majority of populations except Host, Chru and Maga, where, despite fluctuations, prevalence was constantly high (Fig. 3A). Trends can also be observed in the prevalence of

Cestoda gen. spp. Again there are populations with consistently low prevalence (Fig. 3B), but a peak in prevalence can be observed in 2011 for multiple populations (Aroi, Daim and Scad). Aside from a drop in Maga and Chru in 2013, prevalence of *D. gasterostei* remains consistently low (below 50%) in numerous populations, whilst maintaining at high prevalence in a number of others (Fig. 3C). In terms of abundance, *P. filicollis* was rare in most populations with

Table 2. Associations between measures parasite occurrence in individual three-spined sticklebacks on North Uist, and extrinsic (year, population) and intrinsic (length, sex) factors, using GLM analysis

	Population		Year		Year × population		Length		Sex	
	χ^2	<i>P</i>	χ^2	<i>P</i>	χ^2	<i>P</i>	Estimate ± s.e.	<i>P</i>	Estimate ± s.e.	<i>P</i>
Parasite species richness	459.9	***	92.2	***	138.78	***	0.0446 ± 0.0659	***	–	–
<i>G. arcuatus</i> abundance	359.5	***	162.9	***	88.52	***	–	–	–	–
<i>G. arcuatus</i> prevalence	74.69	***	–	–	–	–	0.0653 ± 0.0148	***	–	–
<i>D. gasterostei</i> abundance	524.7	***	75.7	***	178.85	***	0.0700 ± 0.0072	***	–	–
<i>D. gasterostei</i> prevalence	262.41	***	28.69	***	67.46	**	0.0868 ± 0.0156	***	–	–
<i>A. gracilis</i> abundance	270.2	***	84.1	***	123.48	***	0.0636 ± 0.0104	***	–	–
<i>A. gracilis</i> prevalence	177.37	***	30.99	***	62.39	**	0.0788 ± 0.0162	***	–	–
<i>S. solidus</i> abundance	328	***	99.1	***	91.78	***	–	–	0.381 ± 0.131	**
<i>S. solidus</i> prevalence	139.02	***	33.98	***	–	–	–	–	0.639 ± 0.205	**
<i>P. filicollis</i> abundance	565.6	***	–	–	–	–	–0.0494 ± 0.0142	***	–	–
<i>P. filicollis</i> prevalence	148.9	***	–	–	–	–	–	–	–	–
Cestoda gen. spp. abundance	329.5	***	304.6	***	66.86	**	0.0653 ± 0.0184	***	–	–
Cestoda gen. spp. prevalence	54.92	***	37.98	***	–	–	–	–	–	–

N = 1130. ‘Population’ was associated with 13 D.F., ‘year’ with 4 D.F., population × year with 39 D.F., and both ‘sex’ and ‘length’ with 1 D.F. Probability values associated with model: ****P* < 0.001, ***P* ≤ 0.01, **P* ≤ 0.05. ‘Estimate’ refers to (1) the estimated parameter (coefficient) of the effect of length, as given by the GLM and (2) the estimated parameter of the effect of sex, males relative to females.




Fig. 3. Year-to-year variation in prevalence of three parasites of three-spined stickleback in eight populations from North Uist: (a) *P. filicollis*, (b) Cestoda gen. spp. and (c) *D. gasterostei*. For all figure components: Aroi —; Bhar ---; Chru ...; Daim ◆; Host ▲; Maga □; Scad ----; Torm -●-. Only eight populations of the 14 studied are shown, to increase clarity. These eight represent the range of variation in all 14.




Fig. 4. Year-to-year variation in abundance of two parasites of three-spined stickleback in eight populations from North Uist illustrating: (A) consistently low abundance of *P. filicollis* and (B) a rise in *S. solidus* abundance. For all figure components: Aroi —; Bhar ---; Chru ...; Daim ◆; Host ▲; Maga □; Scad ----; Torm -●-. Only eight populations of the 14 studied are shown, to increase clarity. These eight represent the range of variation in all 14.

high counts only observed in Host (Fig. 4A). *Schistocephalus solidus* was rare or absent in many populations, but was consistently present in others (Bhar and Host). It showed a gradual increase across years in Host and a general trend appears to be an increase in later years samples (Fig. 4B).

Variation in mean abundance and prevalence of parasites: (ii) the population level climatic and seasonal effects. Species richness varied greatly between populations (Table 3), as did the prevalence and abundance of all species excluding Cestoda gen. spp. Temperature (in the immediately preceding March and April) had no significant effect on

parasites and their overall species richness. However, species richness increased later in the year (Julian date, Table 3) as did prevalence and abundance of *G. arcuatus*.

Environmental effects. There were few significant relationships between environmental variables and overall average measures of parasite occurrence for lochs (Table 4). Prevalence and abundance of *G. arcuatus* was correlated with *P. pungitius* density. *Schistocephalus solidus* prevalence and both prevalence and abundance of Cestoda gen. spp. were significantly correlated with both calcium concentration and pH. All correlations with calcium concentration

Table 3. Associations between measures of average annual parasite occurrence in three-spined sticklebacks on North Uist, and extrinsic (population, temperature, Julian date) and intrinsic (average length, sex ratio) factors, using GLM analysis for species richness, abundance and prevalence of *G. arcuatus*, *D. gasterostei*, *A. gracilis*, *S. solidus*, *P. filicollis* and Cestoda gen. spp. Sample size = 57 lake + year combinations, based on 1130 fish.

	Population Wald <i>F</i>	<i>P</i>	Temperature <i>P</i>	Julian date Estimate ± s.e.	Wald <i>F</i>	<i>P</i>
Species richness	15.7	***	–	0.023 ± 0.007	10.4	**
log(<i>G. arcuatus</i> abundance + 1)	3.6	***	–	0.017 ± 0.006	7.5	**
<i>G. arcuatus</i> prevalence	6.4	***	–	0.75 ± 0.26	8.4	**
log(<i>D. gasterostei</i> abundance + 1)	8.6	***	–	–	–	–
<i>D. gasterostei</i> prevalence	14.3	***	–	–	–	–
log(<i>A. gracilis</i> abundance + 1)	3.8	***	–	–	–	–
<i>A. gracilis</i> prevalence	6.6	***	–	–	–	–
log(<i>S. solidus</i> abundance + 0.1)	6.8	***	–	–	–	–
<i>S. solidus</i> prevalence	4.6	***	–	–	–	–
log(<i>P. filicollis</i> abundance + 0.1)	14.1	***	–	–	–	–
<i>P. filicollis</i> prevalence	12.7	***	–	–	–	–
Cestoda gen. spp. abundance	–	–	–	–	–	–
Cestoda gen. spp. prevalence	–	–	–	–	–	–

Population is associated with 13 D.F., all other variables are associated with 1 D.F. Probability values associated with model: *** $P < 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$ before correction, significance value α ($P = 0.05$) corrected using sequential Bonferroni correction ($c = 5$). 'Estimate' refers to the estimated parameter of the effect of mean length, temperature and Julian date as given by the GLM.

were positive. Abundance of Cestoda gen. spp. was positively correlated with pH, whereas *S. solidus* and Cestoda gen. spp. prevalence were negative correlated, indicating higher prevalence of these parasites in more acidic lochs. A greater mean abundance of *P. filicollis* was also observed in lochs with higher calcium levels.

DISCUSSION

General (population)

Comparison of the macroparasite communities of three-spined sticklebacks collected from 14 populations across 5 years was used to look for spatio-temporal patterns in parasite occurrence and suggest possible mechanistic explanations behind observed patterns. Whilst variation occurred among populations, in general, infection levels were high: in half of the populations observed, more than 80% of fish examined were infected with at least one parasite and only one population had fewer than 60% of fish infected. Compared with other locations, North Uist sticklebacks exhibit a relatively narrow range of parasite fauna (de Roij and MacColl, 2012): the average species richness in the most diverse loch was 3.7 compared with a mean species richness found to be as high as 5.3 in a study of four localities in the Baltic Sea (Zander, 2007). Despite this, multiple infections were fairly common and in seven of the 14 populations over 50% of fish harboured more than one parasite. The most frequently encountered macroparasites were the monogenean *G. arcuatus*, the trematodes

D. gasterostei and *A. gracilis*, the cestodes *S. solidus* and *P. filicollis* and the Cestoda gen. spp. group, composed of larval *B. scorpii* and *E. crassum*.

Variation in abundance and individual prevalence of parasites: (i) the individual host level. In many previous studies, little evidence was found for repeatability in parasite community patterns across space and/or time (Behnke *et al.* 2008b; Kennedy, 2009), although there are instances which demonstrate some extent of repeatability in measures of parasite community composition (Kennedy, 1993; Carney and Dick, 2000; de Roij and MacColl, 2012). Two long-term studies have investigated parasites communities of eels (*Anguilla Anguilla*) in two English rivers; River Clyst (Kennedy, 1993) and River Otter (Kennedy, 1997). Both studies considered a range of community measures including species composition, richness, dominance and diversity. Considerable and erratic variation was observed between years in both studies, showing a lack of predictability. However, changes in community diversity and dominance in the River Clyst were small, suggesting an underlying stability in community structure. The previous study on North Uist covered 2 years and showed little change in the relative difference in parasite community measures across populations, demonstrating short-term stability in the spatial variation of macroparasite communities (de Roij and MacColl, 2012). This being said, it is important to consider long-term studies in a range of locations before presuming general trends in parasite communities (Kennedy, 1997).

Table 4. Associations between annual averages of measures of parasite occurrence in three-spined sticklebacks on North Uist, and environmental factors, using GLM analysis for species richness, abundance and prevalence of *G. arcuatus*, *D. gasterostei*, *A. gracilis*, *S. solidus*, *P. filicollis* and Cestoda gen. spp.

	Surface area <i>P</i>	Mean depth Estimate ± S.E. <i>P</i>	Mean pH Estimate ± S.E. <i>P</i>	Ca ²⁺ concentration Estimate ± S.E. <i>P</i>	Stickle-back density <i>P</i>	<i>P. pungitius</i> density Estimate ± S.E. <i>P</i>
Log(Species richness)	–	N/A	N/A	0.40 ± 0.15 *	–	–
<i>G. arcuatus</i> abundance	N/A	–	–	–	–	0.25 ± 0.09 *
<i>G. arcuatus</i> prevalence	N/A	–	–	–	–	0.07 ± 0.03 *
<i>D. gasterostei</i> abundance	N/A	–	–	–	–	–
<i>D. gasterostei</i> prevalence	N/A	–	–	–	–	–
<i>A. gracilis</i> abundance	N/A	–	–	–	–	–
<i>A. gracilis</i> prevalence	N/A	–0.10 ± 0.04-	*	–	–	–
log(<i>S. solidus</i> abundance)	N/A	0.70 ± 0.30	***	–	–	–
log(<i>S. solidus</i> prevalence)	N/A	–	–	–2.05 ± 0.59 **	8126 ± 2455 **	–
log(<i>P. filicollis</i> abundance)	N/A	–	–	–	–	–
log(<i>P. filicollis</i> prevalence)	N/A	–	–	–	3776 ± 1409-	–
log(Cestoda gen. spp. abundance + 0.1)	N/A	–	–	1.24 ± 0.39 **	6529 ± 1615 **	–
Cestoda gen. spp. prevalence	N/A	–	–	–0.15 ± 0.05 **	717 ± 194 **	–

Sample size = 14 populations using data from 1130 fish. All variables are associated with 1 D.F. Probability values associated with model: *** $P < 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$ before correction, significance value α ($P = 0.05$) corrected using sequential Bonferroni correction ($c = 7$). ‘Estimate’ refers to the estimated parameter of the effect of pH and calcium concentration, as given by the GLM. Shaded cells indicate comparisons absent from the model.

The present investigation extends the research of de Roij and MacColl (2012) to look for longer-term repeatability, using data from five sampling years, spanning an 8-year period. The present study showed less temporal stability than de Roij and MacColl (2012); however, some measures of parasite community still exhibited substantial consistency across years (prevalence of *G. arcuatus*, *S. solidus*, *P. filicollis* and Cestoda gen. spp. and abundance of *P. filicollis*). The consistency observed in our study indicated that, whilst we were unable to identify clear and predictable patterns in parasite distribution, parasite infections are not stochastic, as concluded in Kennedy (2009). Instead certain parasites are consistently more or less persistent in different locations suggesting that the occurrence of parasites in fish lies somewhere between random and structured communities.

Fish length accounted for some variation in most parasite measures, excluding the abundance of *G. arcuatus* and *S. solidus* and prevalence of *S. solidus*, *P. filicollis* and Cestoda gen. spp. In general, length was positively correlated with measures of parasite infection, apart from *P. filicollis* abundance, with which it was negatively correlated. This is consistent with previous observations regarding the association between length and parasite burden. A comparison of published data comparing length and parasite species richness showed that correlations between them are usually positive (Poulin, 1997). Correlations have also been observed between fish length and intensity of infection with larval digenes and cestodes (Poulin, 2000). There are a number of potential explanations for observed correlations between body length and parasite load. Firstly, the bodies of longer fish have a greater surface area and thus a larger area for parasites to infect (Arneberg *et al.* 1998). Secondly, length is usually associated with the age of fish, so that longer (older) fish have had more time to become infected by parasites and accumulate parasite infections (Behnke *et al.* 2001). This effect of age would be more important in some lochs than others as the age structure within populations varies across North Uist. Many of the lochs contain annual populations, but some lochs are home to individuals living up to 3 years (as observed in Reiv, Maga and Mora, A. R. Singkam, unpublished data). These lochs may therefore contain fish which have accrued parasites over a number of years, possibly resulting in greater burdens in longer, older fish.

The negative correlation observed between fish length and the abundance of *P. filicollis* is supported by early work looking at the seasonal changes in this parasite, which showed that smaller fish exhibited higher infection intensity (Hopkins, 1959). Such variation is suggested to be as a result of different feeding habits based on the observation that smaller stomachs of fish under 1-year-old contained more zooplankton (hosts for *P. filicollis*), whereas

larger fish, older than 1 year, tended to have stomachs containing algae and chironomid larvae, thus are less likely to become infected with *P. filicollis* (Hopkins, 1959). Consideration of the life cycle of *P. filicollis* is consistent with this observation. Once mature, *P. filicollis* migrates to the posterior end of the host intestine in order to release eggs via the anus of the host (Hopkins, 1959). After release of eggs, empty proglottids degenerate, until the entire worm is shed (Meggett, 1914). Field studies on numerous species of *Proteocephalus* have indicated that this maturation and degradation of parasites occurs within a year (Scholz, 1999) after which cestodes are lost from the host. Therefore, if the diet of smaller fish increases their chance of infection, these infections are not persistent enough to be observed in older, larger fish.

The association of parasite communities with sex of fish was less consistent, only explaining variation in abundance and prevalence of *S. solidus*, whereby males were more highly parasitized. This may be explained by mating characteristic of males, both behaviourally and chemically (Folstad *et al.* 1994). Males attract females using bright red colouration, produced by carotenoids, which are acquired via consumption of carotenoid rich foods, such as copepods (Ostlund-Nilsson *et al.* 2010). Copepods are also an important transmitter of a number of stickleback parasites, including *S. solidus* and *P. filicollis*, thus increased secondary sexual colouration also increases exposure to parasites, possibly explaining the higher rate and level of *S. solidus* infection in males (Folstad *et al.* 1994). Furthermore, altered androgen profiles result in immunocompromised males during the breeding season (Folstad and Karter, 1992); thus, sex can affect parasite infection and intensity.

An alternative explanation is that the higher infection observed in males could be a result of sampling bias based on the time of year samples were collected. During the non-breeding seasons, males and females move around in shoals; however, during the mating season breeding males build and defend a nest (Pressley, 1981). Samples were collected using minnow traps set around the borders of the lochs, which will catch only fish found in these areas. As samples were collected during the breeding season, it is likely that many breeding males would have been defending nests at the time, thus samples may be biased towards females and non-breeding males (Bagamian *et al.* 2004). This may also explain, at least in part, the heavily female biased sex ratios observed in data samples. It is also worth noting the hypothesis proposed by Lester (1971) that *S. solidus* infected fish move into shallower waters as a result of oxygen stress, so could have an increased chance of being caught in minnow traps. However, this is unlikely to be a problem in North Uist as lochs are shallow and

movement of water by the wind means the water is well oxygenated throughout (A.D.C. MacColl, personal observations).

Variation in mean abundance and prevalence of parasites: (ii) the population level:

Climatic and seasonal effects. The strong population effects observed for the majority of parasite measures (excluding abundance and prevalence of Cestoda gen. spp.) are consistent with our finding that infection with some parasites differs between populations. Although temperature variation had no effect on the parasites present, the time in the year at which samples were collected did affect the species richness and both prevalence and abundance of *G. arcuatus*, all of which increased when samples were collected at a later point in time between late April and late May. *Gyrodactylus salaris*, a gyrodactylid infecting Atlantic salmon (*Salmo salar*), was observed by Appleby and Mo (1997) to demonstrate seasonal patterns: infection levels were lowest in winter and early spring (following low water temperatures) and increased throughout spring. This is consistent with our findings of greater *G. arcuatus* infection later in the season.

Environmental effects. Nine-spined stickleback density was found to be positively associated with the prevalence and abundance of *G. arcuatus*. This is consistent with previous findings of increased transmission of gyrodactylid species, such as *G. salaris*, which are able to infect both hosts when high densities of both three-spined and nine-spined stickleback are found (Soleng *et al.* 1999). Alternatively, it may be that presence of nine-spined stickleback is indicative of some unmeasured aspects of water chemistry, which are favourable to both *Gyrodactylus* and *P. pungitius* (MacColl *et al.* 2013).

Other environmental variables which correlated with parasite occurrence were pH and calcium concentration. Calcium concentration is commonly found to affect the presence of digenean parasites, for example, Curtis and Rau (1980) found calcium concentration to be associated with *Diplostomum* sp. distribution, as their life cycle requires snail hosts which use calcium for shell production (Cribb *et al.* 2003). Marcogliese and Cone (1996) observed a similar effect of pH on digenes infecting American eels (*Anguilla rostrata*), which were absent from rivers with a pH too low to support their molluscan intermediate host. Therefore it is surprising that the calcium concentration and pH did not explain variation of Digenea between populations. However, calcium concentration was positively correlated with *S. solidus*, *P. filicollis* and Cestoda gen. spp. prevalence, as well as Cestoda gen. spp. abundance. There was little support for these findings in the literature, as calcium

concentration is not commonly found to affect the occurrence of cestodes. We considered the possibility that calcium may be correlated with another variable which could affect the presence of cestodes, perhaps by influencing the presence of copepod intermediate hosts, but this remains an area which will require further study. The pH was positively correlated with the abundance of Cestoda gen. spp. This positive correlation for both calcium concentration and pH observed for this variable is consistent with the findings of MacColl *et al.* (2013). A more surprising result was the negative correlation observed between pH and the prevalence of *S. solidus* and Cestoda gen. spp. *Bothriocephalus claviceps* and *Proteocephalus microcephalus* have both previously been identified in freshwater American eels (*A. rostrata*) living in rivers with pH 4.7–5.0, demonstrating that cestodes are suited to living in harsh aquatic environments (Marcogliese and Cone, 1996). However, these environmental results are puzzling as pH and calcium concentration are usually positively correlated due to dissolved alkaline metals increasing the pH of water (MacColl *et al.* 2013). Thus, one would expect calcium and pH to both be either positively or negatively correlated with parasites, rather than show an inverse relationship. A study by Fryer (1980) found that more acidic lakes were associated with a decreased diversity of crustacean species. It is possible that species able to transmit these cestodes are more suited to survival in acidic lochs than other crustaceans, increasing the chance of sticklebacks consuming infected prey. This idea could be explored with analysis of zooplankton present in lochs.

This study successfully identifies some level of repeatability in parasites infecting North Uist sticklebacks. Although a number of parasites differ in relative abundance and prevalence across the years, consistency was identified with regards to differences between populations in the prevalence of *G. arcuatus*, *S. solidus*, *P. filicollis* and Cestoda gen. spp. and abundance of *P. filicollis* throughout the study, indicating that parasite occurrence is not fully stochastic. Variation in temperature and season had very little effect on parasite distributions but some correlation was identified between parasites and abiotic environmental factors.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182016001815>.

ACKNOWLEDGEMENTS

We are grateful to North Uist Estates and the Scottish Government (SEERAD) for access to land on North Uist. We thank Job de Roij, Aliya El Nagar, Sarah Forbes, Muayad Mahmud, Mark Mainwaring, Shaun Robertson and Jim Whiting for assistance with data

collection. The manuscript was improved by comments from Jerzy Behnke and Andrew Fenton.

FINANCIAL SUPPORT

The work was supported by the U.K. Natural Environment Research Council (Grant no. NE/C517525/1) and the University of Nottingham (both to A. D. C. M.).

REFERENCES

- Abu-Madi, M. A., Behnke, J. M., Lewis, J. W. and Gilbert, F. (1998). Descriptive epidemiology of *Heligmosomoides polygyrus* in *Apodemus sylvaticus* from three contrasting habitats in south-east England. *Journal of Helminthology* **72**, 93–100.
- Andersen, K. and Valtonen, E. (1990). On the infracommunity structure of adult cestodes in freshwater fishes. *Parasitology* **101**, 257–264.
- Anderson, R. M. and May, R. M. (1987). Regulation and stability of host-parasite population interactions: I. Regulatory processes. *Journal of Animal Ecology* **47**, 219–247.
- Appleby, C. and Mo, T. A. (1997). Population dynamics of *Gyrodactylus salaris* (Monogenea) infecting Atlantic salmon, *Salmo salar*, parr in the River Batnfjordseiva, Norway. *Journal of Parasitology* **83**, 23–30.
- Arneberg, P. (2002). Host population density and body mass as determinants of species richness in parasite communities: comparative analyses of directly transmitted nematodes of mammals. *Ecography* **25**, 88–94.
- Arneberg, P., Skorpung, A. and Read, A. F. (1998). Parasite abundance, body size, life histories, and the energetic equivalence rule. *American Naturalist* **151**, 497–513.
- Bagamian, K., Heins, D. and Baker, J. (2004). Body condition and reproductive capacity of three-spined stickleback infected with the cestode *Schistocephalus solidus*. *Journal of Fish Biology* **64**, 1568–1576.
- Behnke, J. (2008). Structure in parasite component communities in wild rodents: predictability, stability, associations and interactions.... or pure randomness? *Parasitology* **135**, 751–766.
- Behnke, J., Bajer, A., Sinski, E. and Wakelin, D. (2001). Interactions involving intestinal nematodes of rodents: experimental and field studies. *Parasitology* **122**, S39–S49.
- Behnke, J., Gilbert, F., Abu-Madi, M. and Lewis, J. (2005). Do the helminth parasites of wood mice interact? *Journal of Animal Ecology* **74**, 982–993.
- Behnke, J., Bajer, A., Harris, P., Newington, L., Pidgeon, E., Rowlands, G., Sheriff, C., Kuliś-Malkowska, K., Siński, E. and Gilbert, F. (2008a). Temporal and between-site variation in helminth communities of bank voles (*Myodes glareolus*) from NE Poland. 1. Regional fauna and component community levels. *Parasitology* **135**, 985–997.
- Behnke, J., Bajer, A., Harris, P., Newington, L., Pidgeon, E., Rowlands, G., Sheriff, C., Kuliś-Malkowska, K., Siński, E. and Gilbert, F. (2008b). Temporal and between-site variation in helminth communities of bank voles (*Myodes glareolus*) from NE Poland. 2. The infracommunity level. *Parasitology* **135**, 999–1018.
- Blanchet, S., Rey, O., Berthier, P., Lek, S. and Loot, G. (2009). Evidence of parasite-mediated disruptive selection on genetic diversity in a wild fish population. *Molecular Ecology* **18**, 1112–1123.
- Bolek, M. G. and Coggins, J. R. (2000). Seasonal occurrence and community structure of helminth parasites from the eastern American toad, *Bufo americanus americanus*, from southeastern Wisconsin, USA. *Comparative Parasitology* **67**, 202–209.
- Bush, A. O., Lafferty, K. D., Lotz, J. M. and Shostak, A. W. (1997). Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* **83**, 575–583.
- Bykhovskaya-Pavloskaya, I. E., Gusev, A. V., Dubinina, M. N., Izyumova, N. A., Smirnova, T. S., Sokolovskaya, I. L., Shtein, G. A., Shul'man, S. S. and Epshtein, V. M. (1964). *Key to Parasites of Freshwater Fish in the U.S.S.R.* Israel Program for Scientific Translations, Jerusalem, Israel.
- Calvete, C., Blanco-Aguilar, J., Virgós, E., Cabezas-Díaz, S. and Villafuerte, R. (2004). Spatial variation in helminth community structure in the red-legged partridge (*Alectoris rufa* L.): effects of definitive host density. *Parasitology* **129**, 101–113.
- Cardon, M., Loot, G., Grenouillet, G. and Blanchet, S. (2011). Host characteristics and environmental factors differentially drive the burden and pathogenicity of an ectoparasite: a multilevel causal analysis. *Journal of Animal Ecology* **80**, 657–667.
- Carney, J. and Dick, T. (2000). Helminth communities of yellow perch (*Perca flavescens* (Mitchill)): determinants of pattern. *Canadian Journal of Zoology* **78**, 538–555.
- Chappell, L. (1969). Competitive exclusion between two intestinal parasites of the three-spined stickleback, *Gasterosteus aculeatus* L. *Journal of Parasitology* **55**, 775–778.
- Connor, E. F. and McCoy, E. D. (1979). The statistics and biology of the species-area relationship. *American Naturalist* **113**, 791–833.
- Cribb, T. H., Bray, R. A., Olson, P. D., Timothy, D. and Littlewood, J. (2003). Life cycle evolution in the Digenea: a new perspective from phylogeny. *Advances in Parasitology* **54**, 197–254.
- Curtis, M. A. and Rau, M. E. (1980). The geographical distribution of diplostomiasis (Trematoda: Strigeidae) in fishes from northern Quebec, Canada, in relation to the calcium ion concentrations of lakes. *Canadian Journal of Zoology* **58**, 1390–1394.
- Dartnall, H. J. G. (1973). Parasites of the nine-spined stickleback *Pungitius pungitius* (L.). *Journal of Fish Biology* **5**, 505–509.
- de Roij, J. and MacColl, A. D. C. (2012). Consistent differences in macroparasite community composition among populations of three-spined sticklebacks, *Gasterosteus aculeatus* L. *Parasitology* **139**, 1478–1491.
- Ebert, D., Hottinger, J. W. and Pajunen, V. I. (2001). Temporal and spatial dynamics of parasite richness in a *Daphnia* metapopulation. *Ecology* **82**, 3417–3434.
- Folstad, I. and Karter, A. J. (1992). Parasites, bright males, and the immunocompetence handicap. *American Naturalist* **139**, 603–622.
- Folstad, I., Hope, A. M., Karter, A. and Skorpung, A. (1994). Sexually selected color in male sticklebacks: a signal of both parasite exposure and parasite resistance? *Oikos* **69**, 511–515.
- Fryer, G. (1980). Acidity and species diversity in freshwater crustacean faunas. *Freshwater Biology* **10**, 41–45.
- Goater, C., Baldwin, R. and Scrimgeour, G. (2005). Physico-chemical determinants of helminth component community structure in whitefish (*Coregonus clupeaformis*) from adjacent lakes in Northern Alberta, Canada. *Parasitology* **131**, 713–722.
- González, M. and Poulin, R. (2005). Spatial and temporal predictability of the parasite community structure of a benthic marine fish along its distributional range. *International Journal for Parasitology* **35**, 1369–1377.
- Halmetoja, A., Valtonen, E. T. and Koskenniemi, E. (2000). Perch (*Perca fluviatilis* L.) parasites reflect ecosystem conditions: a comparison of a natural lake and two acidic reservoirs in Finland. *International Journal for Parasitology* **30**, 1437–1444.
- Hopkins, C. (1959). Seasonal variations in the incidence and development of the cestode *Proteocephalus filicollis* (Rud. 1810) in *Gasterosteus aculeatus* (L. 1766). *Parasitology* **49**, 529–542.
- Kennedy, C. R. (1993). The dynamics of intestinal helminth communities in eels *Anguilla anguilla* in a small stream: long-term changes in richness and structure. *Parasitology* **107**, 71–78.
- Kennedy, C. R. (1997). Long-term and seasonal changes in composition and richness of intestinal helminth communities in eels *Anguilla anguilla* of an isolated English river. *Folia Parasitologica* **44**, 267–273.
- Kennedy, C. R. (2009). The ecology of parasites of freshwater fishes: the search for patterns. *Parasitology* **136**, 1653–1662.
- Kennedy, C. R., Shears, P. and Shears, J. (2001). Long-term dynamics of *Ligula intestinalis* and roach *Rutilus rutilus*: a study of three epizootic cycles over thirty-one years. *Parasitology* **123**, 257–269.
- Lawton, J. H. (1999). Are there general laws in ecology? *Oikos* **84**, 177–192.
- Lester, R. (1971). The influence of *Schistocephalus plerocercoids* on the respiration of *Gasterosteus* and a possible resulting effect on the behavior of the fish. *Canadian Journal of Zoology* **49**, 361–366.
- Lively, C. M., de Roode, J. C., Duffy, M. A., Graham, A. L. and Koskella, B. (2014). Interesting open questions in disease ecology and evolution. *American Naturalist* **184**, S1–S8.
- MacColl, A. D. C. (2009). Parasite burdens differ between sympatric three-spined sticklebacks. *Ecography* **32**, 153–160.
- MacColl, A. D. C., Nagar, A. E. and de Roij, J. (2013). The evolutionary ecology of dwarfism in three-spined sticklebacks. *Journal of Animal Ecology* **82**, 642–652.
- Magurran, A. E. (2004). *Measuring Biological Diversity*, 2nd Edn. Blackwell Science Ltd, Oxford, UK.
- Marcogliese, D. J. and Cone, D. K. (1991). Importance of lake characteristics in structuring parasite communities of salmonids from insular Newfoundland. *Canadian Journal of Zoology* **69**, 2962–2967.
- Marcogliese, D. J. and Cone, D. K. (1996). On the distribution and abundance of eel parasites in Nova Scotia: influence of pH. *Journal of Parasitology* **82**, 389–399.

- Meggitt, F. J.** (1914). The structure and life history of a tapeworm (*Ichthyotaenia filicollis* Rud.) parasitic in the stickleback. *Proceedings of the Zoological Society of London* **1**, 113–138.
- Ostlund-Nilsson, S., Mayer, I. and Huntingford, F. A.** (2010). *Biology of the Three-spined Stickleback*. CRC Press: Taylor & Francis Group, Florida, USA.
- Pennycuik, L.** (1971). Seasonal variations in the parasite infections in a population of three-spined sticklebacks, *Gasterosteus aculeatus* L. *Parasitology* **63**, 373–388.
- Pietroock, M. and Marcogliese, D. J.** (2003). Free-living endohelminth stages: at the mercy of environmental conditions. *Trends in Parasitology* **19**, 293–299.
- Poulin, R.** (1997). Species richness of parasite assemblages: evolution and patterns. *Annual Review of Ecology and Systematics* **28**, 341–358.
- Poulin, R.** (2000). Variation in the intraspecific relationship between fish length and intensity of parasitic infection: biological and statistical causes. *Journal of Fish Biology* **56**, 123–137.
- Poulin, R.** (2007). Are there general laws in parasite ecology? *Parasitology* **134**, 763–776.
- Pressley, P. H.** (1981). Parental effort and the evolution of nest-guarding tactics in the three-spined stickleback *Gasterosteus aculeatus* L. *Evolution* **35**, 282–295.
- Scholz, T.** (1999). Life cycles of species of *Proteocephalus* parasites of fishes in the Palearctic Region: a review. *Journal of Helminthology* **73**, 1–19.
- Soleng, A., Jansen, P. A. and Bakke, T. A.** (1999). Transmission of the monogenean *Gyrodactylus salaris*. *Folia Parasitologica* **46**, 179–184.
- Vidal-Martínez, V. and Poulin, R.** (2003). Spatial and temporal repeatability in parasite community structure of tropical fish hosts. *Parasitology* **127**, 387–398.
- Waterston, A. R., Holden, A. V., Campbell, R. N. and Maitland, P. S.** (1979). The inland waters of the Outer Hebrides. *Proceedings of the Royal Society of Edinburgh Section B: Biological Sciences* **77B**, 329–351.
- Wilson, K., Bjørnstad, O., Dobson, A., Merler, S., Pogliayen, G., Randolph, S., Read, A. and Skorping, A.** (2002). Heterogeneities in macroparasite infections: patterns and processes. *The Ecology of Wildlife Diseases* **44**, 6–44.
- Zander, C. D.** (2007). Parasite diversity of sticklebacks from the Baltic Sea. *Parasitology Research* **100**, 287–297.