

Resolving variation in the reproductive tradeoff between sperm size and number

Simone Immler^{a,b,1,2}, Scott Pitnick^c, Geoff A. Parker^d, Kate L. Durrant^c, Stefan Lüpold^c, Sara Calhim^e, and Tim R. Birkhead^a

^aDepartment of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield S10 2TN, United Kingdom; ^bDepartment of Evolutionary Biology, Evolutionary Biology Centre, Uppsala University, 75236 Uppsala, Sweden; ^cDepartment of Biology, Syracuse University, Syracuse, NY 13244; ^dDepartment of Evolution, Ecology and Behaviour, Institute of Integrative Biology, University of Liverpool, Liverpool L69 7ZB, United Kingdom; and ^eDepartment of Biology, Queen's University, Kingston, ON, Canada K7L 3N6

Edited by Nina Wedell, University of Exeter, Exeter, Devon, United Kingdom, and accepted by the Editorial Board February 14, 2011 (received for review June 30, 2010)

Spermatozoa are amongst the most variable cells, and three factors are thought to account for this variation in design: fertilization mode, phylogeny, and postcopulatory sexual selection. In addition, it has long been assumed that a tradeoff exists between sperm size and number, and although postcopulatory sexual selection affects both traits, empirical evidence for a tradeoff has so far been elusive. Our recent theoretical model predicts that the nature of a direct tradeoff between sperm size and number varies with sperm competition mechanism and sperm competition risk. We test these predictions using a comparative approach in two very different taxa with different sperm competition mechanisms: passerine birds (mechanism: simple raffle) and *Drosophila* fruit flies (sperm displacement). We show that in both groups, males increase their total ejaculate investment with increasing sperm competition risk, but whereas passerine birds allocate disproportionately to sperm number, drosophilids allocate disproportionately to sperm size. This striking difference between the two groups can be at least partly explained by sperm competition mechanisms depending on sperm size relative to the size of the female reproductive tract: in large animals (passerines), sperm numbers are advantageous in sperm competition owing to dilution inside the female tract, whereas in small animals (drosophilids), large sperm are advantageous for physical competition (sperm displacement). Our study provides two important results. First, we provide convincing evidence for the existence of a sperm size–number tradeoff. Second, we show that by considering both sperm competition mechanism and dilution, we can account for variation in sperm size between different taxa.

dilution effect | sperm allocation

The seemingly unlimited number of tiny sperm produced by males compared with the few large eggs produced by females (i.e., anisogamy) has shaped our understanding of sexual selection and the evolution of mating systems (1–3). However, although the males of most animals produce huge numbers of tiny sperm, some produce few, giant sperm (4), suggesting that males trade off sperm size and sperm number. Theoretical models of sperm size evolution in the context of sperm competition assume either a direct or indirect tradeoff between sperm size and sperm number. Under a direct tradeoff (5), males have a fixed energy budget, M , to invest in an ejaculate, so if each sperm has size m , the sperm number will be $s = M/m$; any change in one strategic parameter (m or s) trades off directly against the other. Under an indirect tradeoff (6), a fixed resource budget, R , is allocated between the ejaculate and effort spent acquiring a mating, C , where the number of matings, n , obtained by a male is given by $R/(C + Dsm)$ and where Dsm defines the amount of energy invested into the ejaculate. Thus, a male can effectively now vary three strategic parameters: n , s , and m . Our recent theoretical model (7) shows that at a given level of R and sperm competition risk q (i.e., at a given level β ; *Box 1*), a direct tradeoff between sperm size and number effectively still operates even under the indirect tradeoff model. However, empirical evidence for such a tradeoff has been equivocal (8–11). The present study couples data with theoretical predictions of our recent model (7) (*Box 1*)

to generate insights into sperm size/number evolution. It introduces the concepts of the mechanisms of sperm competition and dilution to explain observed interspecific variation in sperm size. Only by considering both of these factors are we able to account for the relationship between body mass and sperm size, as well as for sperm gigantism in *Drosophila*.

To clarify the logic, we give an overview of our recent theoretical model of sperm size–number tradeoffs and its main predictions in *Box 1*; the full account is complex and is presented elsewhere (7). It differs from previous models (6) in predicting how absolute and relative expenditure on sperm size and number should vary in relation to sperm competition risk, q , in both sperm mixing and sperm displacement systems, under varying sperm density: (i) sperm number, s , should always increase with q , but sperm size (defined here as mass), m , may be constant, increasing, or decreasing with q , (ii) total ejaculate investment, ms , should always increase with q , and (iii) the ratio m/s (which indicates how allocation within ms is skewed toward size or number as q increases) can increase or decrease with q depending on the influence of sperm density (the density of sperm in the set from which fertilizations are drawn) on sperm competition; specifically, (iv) if sperm size has no effect (or only a moderate effect) on sperm competitive ability with increasing sperm density (e.g., in systems with a raffle mechanism, such as external fertilizers and internally fertilizing vertebrates), then sperm number is favored over sperm size with increasing risk of sperm competition (thus as ms increases with q , the ratio m/s decreases), or (v) if sperm competitive ability is strongly affected by increases in sperm density (e.g., where sperm displacement is an important mechanism), then sperm size is favored over sperm number with increasing risk of sperm competition (thus as ms increases with q , m/s may also increase). Here we examine the above predictions in two taxonomic groups with internal fertilization, which differ considerably in their sperm competition mechanisms and sperm densities: passerine birds, in which sperm competition obeys a simple raffle mechanism (12), and *Drosophila* fruit flies, in which sperm competition success relies on displacement mechanisms (13, 14).

To examine relationships between sperm size and number and sperm competition across species, we use relative testis size as an index of sperm competition risk (a causal positive correlation between the two has been demonstrated in numerous experimental and comparative studies) (15–23). Relative testis size is also typically a direct reflection of male investment in sperm production (24, 25) and therefore likely to correlate with the cost

Author contributions: S.I. and T.R.B. designed research; S.I., S.P., G.A.P., K.L.D., S.L., and S.C. performed research; S.I. analyzed data; and S.I., S.P., G.A.P., and T.R.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. N.W. is a guest editor invited by the Editorial Board.

¹Present address: Evolutionary Biology Center/Evolutionary Biology, University of Uppsala, 752 36 Uppsala, Sweden.

²To whom correspondence should be addressed. E-mail: simone.immler@ebc.uu.se.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1009059108/-DCSupplemental.

of each ejaculate, proportional to the product of the evolutionarily stable strategy (ESS) sperm size and number in our model (i.e., m^*s^* : as expected, our data demonstrate such a positive relationship in both birds and flies) (see below). Our focus here, however, is on how the tradeoff balance between sperm size and number changes with increasing sperm competition risk (and hence increasing m^*s^*) and how this balance is affected by systems with different sperm competition mechanisms and sperm densities (see below). In particular, from the predictions above, we expect passerines, with their low sperm density raffle mechanism, to show a decreasing relationship between m^*s^* and risk q (Box 1 and Fig. 1C). In sharp contrast, the drosophilids, with their high-density sperm displacement mechanism, are predicted to show an increasing relationship between m^*s^* and risk q (Box 1 and Fig. 1D).

We next present a hypothesis that arises (a priori) from our theoretical models, basic biology, and empirical data on sperm competition in different groups; these collectively enable strong quantitative predictions that can be compared with patterns within animal groups. Our rationale is as follows. Vertebrate systems with internal fertilization typically have low sperm densities around the egg during fertilization, owing to reproductive tracts that are large relative to sperm size (26–28). Consequently, competitive advantages of increased sperm size in the region of the ESS are likely to be relatively weak across wide ranges of natural sperm densities and related mainly to effects of size on

sperm motility and survival. Further, in birds, the sperm competition mechanism obeys the raffle principle (12, 29), which requires strongly escalating competitive benefits of sperm size with sperm density to generate disproportionate allocation to sperm size with increased sperm competition risk. It is difficult to envisage that passerines could fulfill these requirements, and so we predict that m^*s^* should decrease with q (Box 1 and Fig. 1C). The same applies to externally fertilizing organisms. In contrast, in insects and other invertebrates with internal fertilization, longer (or larger) sperm have been shown to have a competitive fertilization advantage against shorter (or smaller) sperm of rival males through enhanced displacement or other competitive abilities (13, 20, 30, 31). In insects, sperm are stored at high density in the female sperm-storage organs (13, 32), which in *Drosophila* have fixed volume so that new ejaculates displace previously stored sperm. At fertilization, sperm compete at high density around the proximal region of the sperm store duct to enter the micropyle of the egg as it passes down the oviduct. Longer sperm seem to provide a competitive advantage mediated through location rather than swimming speed (13). For this scenario, our models predict sperm size to increase more than sperm number with increasing risk of sperm competition (m^*s^* increasing; Box 1 and Fig. 1D) only under special circumstances, most easily met in displacement systems, when sperm size has special and strong competitive advantages in skewing fertilization, notably if the minimum successful sperm size escalates as sperm density increases (7, 13). The propensity for increasingly escalating competitive benefits of sperm size with sperm density in systems such as *Drosophila* is considerable, and some of the largest sperm described to date are found in small-bodied invertebrate taxa (33, 34).

The well-known phenomenon of sperm density declining as the female reproductive tracts become larger (the “sperm dilution” effect) (15, 35–37) may act in a manner that reinforces our predictions. In large animals, the dilution effect hypothesis suggests a compensating increase in sperm number at the expense of sperm size—and hence female tract dimensions—increases (assuming that female reproductive tract size scales directly with body size). In tiny animals, by contrast, dilution constraints on the evolution of longer sperm are largely absent because sperm size is relatively large compared with body size (33), and furthermore, sperm are typically stored in small, fixed-volume storage organs.

Using a comparative approach, we tested our predictions by quantifying the relationship between sperm size, sperm number, and the total investment in spermatogenesis (m^*s^*) in passerines (Aves, Passeriformes) as a relatively large-bodied taxon [5.7–589 g body mass (BM)] obeying the raffle principle, and in *Drosophila* (Insecta, Diptera) as a small-bodied taxon (176–1,020 mg BM) obeying sperm displacement. These two taxa certainly differ in many biological aspects, which are not taken into account here, and we caution that a comparative analysis of two lineages cannot establish causation. However, because sperm competition mechanisms are well understood in both taxa, we believe that the differences between sperm size and number in the two groups are likely to be at least partly determined by the selective forces outlined above. Our results generally supported our theoretical predictions and provide intriguing information about mechanisms involved in postcopulatory sexual selection and animal reproduction in general.

Results

For passerines, we collected data on sperm size, combined testis mass (CTM), and BM for a total of 196 species (Table S1), and sperm number for 23 of these species. As a proxy for ejaculate size we used the number of sperm counted in the seminal glomera (i.e., the extragonadal male sperm-storage organs). Sperm numbers in the seminal glomera are directly related to testis size (17), which in turn is related to ejaculate size (16). Although the use of this proxy may add random noise due to measurement errors, it will not bias our results in a systematic way and hence does not jeopardize the overall conclusions we can draw from




Fig. 1. Theoretical predictions for sperm size–number relationship with increasing risk of sperm competition. (A) The ESS total sperm expenditure (product of sperm size and number, m^*s^*) always increases across the risk range sperm competition risk, q , as observed in passerine birds and *Drosophila* (Fig. 3 A and B). (B) Sperm loading factor, r , determines a sperm's competitive ability and must increase at a decreasing rate with m . The tangent to the curve $r(m)$ gives the marginal value theorem solution for ESS sperm size, m^* , at a given level of sperm competition risk, q . If sperm density increases the intercept of $r(m)$, or decreases its slope, a higher ESS sperm size is generated. (C) The “default expectation” (as observed in passerine birds; Fig. 3C)—a decreasing relationship between the ratio of sperm size/number, m^*/s^* , and sperm competition risk, q , is expected when (i) $r(m)$ is unaffected by the sperm density, or where (ii) increasing sperm density decreases the intercept of $r(m)$ or increases the slope of $r(m)$, or where (iii) increasing sperm density only generates weak increases in the intercept of $r(m)$ or weak decreases in the slope of $r(m)$. (D) An increasing relationship between the ratio of sperm size/number, m^*/s^* , and sperm competition risk, q (as observed in *Drosophila*; Fig. 3D), is expected only when increasing sperm density sharply decreases the slope or increases intercept of $r(m)$ and is more likely to occur in sperm displacement systems such as those found in insects.




Fig. 3. Sperm size–number relationship with increasing risk of sperm competition. In both (A) passerine birds and (B) *Drosophila*, the overall investment in sperm size (m^*) and number (s^*) increases with increasing risk of sperm competition (compare with theoretical prediction in Fig. 1A). As predicted (Box 1) (C; compare with Fig. 1C), in passerine birds sperm number increases at the expense of sperm size, whereas (D; compare with Fig. 1D) in *Drosophila* sperm size increases at the expense of sperm number. Figures are not controlled for phylogeny.

suggest that this phenomenon will be mainly restricted to displacement systems in which sperm size plays an increasingly crucial role in sperm competition as sperm density increases.

Thus, in support of our theoretical predictions (Box 1): (i) total ejaculate investment, m^*s^* , increases with relative testis size (sperm competition risk) in both passerine birds and fruit flies, (ii) relative investment in sperm size, m^*/s^* , decreases with sperm competition risk in passerines but increases in *Drosophila*, and (iii) sperm numbers, s^* , increase with risk in passerines and also across the low-risk range in *Drosophila*. Only the decline in sperm numbers across the high-risk range in *Drosophila* fails to concur with our expectations.

Discussion

Our findings suggest that males are forced to trade off sperm size against sperm number and that this tradeoff may go either way. These results have important implications for our understanding of sexual selection and the evolution of mating systems, because the difference in size and number between male and female gametes determines to a large extent the strength of sexual selection (20). In addition, our evidence for a tradeoff between sperm size and number supports the idea that sperm production is costly and that males are under pressure to adaptively allocate their reproductive energy, particularly under high risk levels of sperm competition. Our results are compatible with our theoretical predictions of a tradeoff between sperm size and number (7) (Box 1) and that as sperm competition risk increases, the balance between sperm size and number can shift in either direction. We postulate that the difference we find between passerine birds and *Drosophila* is based on the sperm competition mechanism, which in turn depends on sperm density and hence on body size and the volume of the female reproductive tract.

In the large-bodied taxon (passerine birds), the advantages of sperm size with increasing sperm density are unlikely to accelerate sufficiently strongly to favor sperm size rather than sperm numbers, and hence sperm number increases more rapidly than (or instead of) sperm size (the “default” expectation in Scenario 1 in our model; Fig. 1C). In contrast, the fact that sperm size increases

disproportionately to sperm numbers in *Drosophila* supports the notion that in this taxon sperm size can play a major role in sperm competition, owing to displacement and direct competition between sperm at the site of fertilization (according to Scenario 2 in our model) (12, 13). Sperm in such species are commonly longer than any distance they travel within the female reproductive tract (19, 41). Moreover, experimental evolution studies of *Drosophila* have demonstrated a positive association between sperm length and postcopulatory sexual selection intensity (19) and revealed that longer sperm are better at displacing shorter sperm—and resisting being displaced by them—from the site of fertilization (12, 13). These findings provide an intriguing explanation for the evolution of “giant” sperm observed in some *Drosophila* species (4, 20). However, we recognize that factors other than sperm competition mechanisms, such as physiological differences (e.g., poikilotherm vs. ectotherm), may certainly also influence the evolution of sperm size and number and contribute to the variation in the evolution of sperm size and number across taxa.

In addition, our study provides a plausible explanation for some of the seemingly contradictory results of previous studies, which have shown no relationship (41, 42), a positive relationship (43–45), or a negative relationship between sperm size and sperm competition (44, 46). This variation across studies can be explained in terms of how the competitive benefits of sperm size, $r(m)$ (Fig. 1B), change with sperm density (Box 1). The differential impact of $r(m)$ may explain the previous findings of a positive relationship between sperm size and risk of sperm competition in the family of Fringillidae and a negative relationship in the Sylviidae (44) (Fig. 2A). The theoretical predictions for the raffle mechanism are that species with an overall low risk of sperm competition q will show a sharply increasing positive relationship between sperm size and q , whereas species with high q will exhibit little or no relation between sperm size and q , because the predicted relation asymptotes quickly with q (7). Our data support these predictions in two ways. First, comparing two families: the Fringillidae have relatively low levels of sperm competition and show a positive relationship between sperm size and q , whereas the Sylviidae have a higher level of sperm com-

testes and BM were obtained for 10 species from Pitnick (24); values for eight further species were obtained using methods described in Pitnick (24).

Statistical Analyses. To account for the statistical nonindependence of data points due to shared ancestry we used a generalized least-squares (GLS) approach in a phylogenetic framework (PGLS; 38–40) to perform multiple regression analyses. The phylogenetic GLS is based on the use of maximum likelihood (ML) models and takes phylogeny into account by referring to an internal matrix of expected covariances among species based on their degree of shared ancestry. In addition, the ML approach allows the estimation of the phylogenetic dependence parameter λ , which ranges between 0 and 1, indicating the relative importance of phylogeny in explaining the similarities between traits. Values of λ close to 0 indicate that the underlying phylogeny explains little of the observed trait variation, whereas values of λ close to 1 indicate strong phylogenetic association of the traits. Analyses were performed using a code developed by Freckleton for the statistical package R V.2.10.1 (53). The phylogenetic topology for passerine birds was

inferred from published sources (Fig. S1), and the phylogenetic topology for *Drosophila* species was adopted from two published sources (54, 55) (Fig. S2). We included sperm traits (size or number) as response variables and in all models testis size and body size as independent variables. We also added a quadratic term for testis size and an interaction term between testis size and BM into the model; however, these were removed if not significant. To optimize the fit of the model, data were transformed accordingly using \ln or square-root transformations. For comparisons of relationships with a slope of 1, we calculated the RMA slope and performed a one-sided t test on the RMA slope compared with 1.

ACKNOWLEDGMENTS. We thank Leigh Simmons for stimulating discussions of this study and Göran Arnqvist and Ted Morrow for comments on earlier drafts of the manuscript. This work was supported by Leverhulme Grant F/00118AJ (to T.R.B.) and National Science Foundation Grants DEB-0814732 and DEB-6990357 (to S.P.).

- Bateman AJ (1948) Intra-sexual selection in *Drosophila*. *Heredity* 2:349–368.
- Trivers RL (1972) Parental investment and sexual selection. *Sexual Selection and the Descent of Man 1871–1971*, ed Campbell B (Aldine-Atherton, Chicago), pp 136–179.
- Arnold SJ (1994) Bateman's principals and the measurement of sexual selection in plants and animals. *Am Nat* 144:s126–s149.
- Pitnick S, Spicer GS, Markow TA (1995) How long is a giant sperm? *Nature* 375:109.
- Parker GA (1982) Why are there so many tiny sperm? Sperm competition and the maintenance of two sexes. *J Theor Biol* 96:281–294.
- Parker GA (1993) Sperm competition games: Sperm size and sperm number under adult control. *Proc Biol Sci* 253:245–254.
- Parker GA, Immler S, Pitnick S, Birkhead TR (2010) Sperm competition games: Sperm size (mass) and number under raffle and displacement, and the evolution of P2. *J Theor Biol* 264:1003–1023.
- Gage MJG, Cook PA (1994) Sperm size or numbers—effects of nutritional stress upon eupyrene and apyrene sperm production strategies in the moth *Plodia interpunctella* (Lepidoptera, Pyralidae). *Funct Ecol* 8:594–599.
- Arnaud L, Haubruge E, Gage MJG (2001) Sperm size and number variation in the red flour beetle. *Zool J Linn Soc* 133:369–375.
- Snook RR (2005) Sperm in competition: Not playing by the numbers. *Trends Ecol Evol* 20:46–53.
- Lessells CM, Snook RR, Hosken DJ (2009) The evolutionary maintenance of sperm: Selection for a small, motile gamete mating type. *Sperm Biology: An Evolutionary Perspective*, eds Birkhead TR, Hosken DJ, Pitnick S (Academic Press, Oxford), pp 43–67.
- Colegrave N, Birkhead TR, Lessells CM (1995) Sperm precedence in zebra finches does not require special mechanisms of sperm competition. *Proc Biol Sci* 259:223–228.
- Pattarini JM, Starmer WT, Bjork A, Pitnick S (2006) Mechanisms underlying the sperm quality advantage in sperm competition and cryptic female choice in *Drosophila melanogaster*. *Evolution* 60:2064–2080.
- Manier MK, et al. (2010) Resolving mechanisms of competitive fertilization success in *Drosophila melanogaster*. *Science* 328:354–357.
- Harcourt AH, Harvey PH, Larson SG, Short RV (1981) Testis weight, body weight and breeding system in primates. *Nature* 293:55–57.
- Møller AP (1988) Testes size, ejaculate quality and sperm competition in birds. *Biol J Linn Soc Lond* 33:273–283.
- Birkhead TR, Briskie JV, Møller AP (1993) Male sperm reserves and copulation frequency in birds. *Behav Ecol Sociobiol* 32:85–93.
- Byrne PG, Roberts JD, Simmons LW (2002) Sperm competition selects for increased testes mass in Australian frogs. *J Evol Biol* 15:347–355.
- Miller GT, Pitnick S (2002) Sperm-female coevolution in *Drosophila*. *Science* 298:1230–1233.
- Bjork A, Pitnick S (2006) Intensity of sexual selection along the anisogamy-isogamy continuum. *Nature* 441:742–745.
- Pitnick S, Miller GT, Reagan J, Holland B (2001) Males' evolutionary responses to experimental removal of sexual selection. *Proc Biol Sci* 268:1071–1080.
- Ramm SA, Parker GA, Stockley P (2005) Sperm competition and the evolution of male reproductive anatomy in rodents. *Proc Biol Sci* 272:949–955.
- Calhim S, Birkhead TR (2007) Testes size in birds: Quality versus quantity—assumptions, errors, and estimates. *Behav Ecol* 18:271–275.
- Pitnick S (1996) Investment in testes and the cost of making long sperm in *Drosophila*. *Am Nat* 148:57–80.
- Schärer L, Vizoso D (2007) Phenotypic plasticity in sperm production rate: There's more to it than testis size. *Evol Ecol* 21:295–306.
- Brillard JP, Bakst MR (1990) Quantification of spermatozoa in the sperm-storage tubules of turkey hens and the relation to sperm numbers in the perivitelline layer of eggs. *Biol Reprod* 43:271–275.
- Suarez SS, Pacey AA (2006) Sperm transport in the female reproductive tract. *Hum Reprod Update* 12:23–37.
- Holt WV (2009) Is semen analysis useful to predict the odds that the sperm will meet the egg? *Reprod Domest Anim* 44 (Suppl 3):31–38.
- Birkhead TR (1998) Sperm competition in birds: mechanisms and function. *Sperm Competition and Sexual Selection*, eds Birkhead TR, Møller AP (Academic Press, London), pp 579–622.
- Radwan J (1996) Intraspecific variation in sperm competition success in the bulb mite: A role for sperm size. *Proc Biol Sci* 263:855–859.
- LaMunyon CW, Ward S (1999) Evolution of sperm size in nematodes: Sperm competition favours larger sperm. *Proc Biol Sci* 266:263–267.
- Sbilordo S, Schäfer M, Ward P (2009) Sperm release and use at fertilization by yellow dung fly females (*Scathophaga stercoraria*). *Biol J Linn Soc Lond* 98:511–518.
- Pitnick S, Hosken DJ, Birkhead TR (2009) Sperm morphological diversity. *Sperm Biology: An Evolutionary Perspective*, eds Birkhead TR, Hosken DJ, Pitnick S (Academic Press, Oxford), pp 69–149.
- Hosken DJ (2003) Dispatch. Sperm biology: Size indeed matters. *Curr Biol* 13:R355–R356.
- Short RV (1981) Sexual selection in man and the great apes. *Reproductive Biology of the Great Apes*, ed Graham CE (Academic Press, New York).
- Cummins JM, Woodall PF (1985) On mammalian sperm dimensions. *J Reprod Fertil* 75:153–175.
- Short R (1979) Sexual selection and its component parts, somatic and genital selection, as illustrated by man and the great apes. *Adv Stud Behav* 9:131–158.
- Pagel M (1999) Inferring the historical patterns of biological evolution. *Nature* 401:877–884.
- Pagel M (1997) Inferring evolutionary processes from phylogenies. *Zool Scr* 26:331–348.
- Freckleton RP, Harvey PH, Pagel M (2002) Phylogenetic analysis and comparative data: A test and review of evidence. *Am Nat* 160:712–726.
- Johnson DDP, Briskie JV (1999) Sperm competition and sperm length in shorebirds. *Condor* 101:848–854.
- Gage MJG, Freckleton RP (2003) Relative testis size and sperm morphology across mammals: No evidence for an association between sperm competition and sperm length. *Proc Biol Sci* 270:625–632.
- Byrne PG, Simmons LW, Roberts JD (2003) Sperm competition and the evolution of gamete morphology in frogs. *Proc Biol Sci* 270:2079–2086.
- Immler S, Birkhead TR (2007) Sperm competition and sperm midpiece size: No consistent pattern in passerine birds. *Proc Biol Sci* 274:561–568.
- Fitzpatrick JL, et al. (2009) Female promiscuity promotes the evolution of faster sperm in cichlid fishes. *Proc Natl Acad Sci USA* 106:1128–1132.
- Stockley P, Gage MJG, Parker GA, Møller AP (1997) Sperm competition in fishes: The evolution of testis size and ejaculate characteristics. *Am Nat* 149:933–954.
- Parker GA, Ball MA (2005) Sperm competition, mating rate and the evolution of testis and ejaculate sizes: A population model. *Biol Lett* 1:235–238.
- Immler S, Birkhead TR (2005) A non-invasive method for obtaining spermatozoa from birds. *Ibis* 147:827–830.
- Wolfson A (1952) The cloacal protuberance—a means for determining breeding condition in live male passerine birds. *Bird-Banding* 23:159–165.
- Pitnick S, Markow TA, Spicer GS (1995) Delayed male maturity is a cost of producing large sperm in *Drosophila*. *Proc Natl Acad Sci USA* 92:10614–10618.
- Schärer L, Da Lage JL, Joly D (2008) Evolution of testicular architecture in the Drosophilidae: A role for sperm length. *BMC Evol Biol* 8:143.
- Dunning JB (1993) *CRC Handbook of Avian Body Masses* (CRC Press, Boca Raton, FL).
- R Development Core Team (2008) *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna).
- Russo CA, Takezaki N, Nei M (1995) Molecular phylogeny and divergence times of drosophilid species. *Mol Biol Evol* 12:391–404.
- Pitnick S, Markow TA, Spicer GS (1999) Evolution of multiple kinds of female sperm-storage organs in *Drosophila*. *Evolution* 53:1804–1822.
- Charnov EL (1976) Optimal foraging, the marginal value theorem. *Theor Popul Biol* 9:129–136.

Corrections

EVOLUTION

Correction for “Resolving variation in the reproductive tradeoff between sperm size and number,” by Simone Immler, Scott Pitnick, Geoff A. Parker, Kate L. Durrant, Stefan Lüpold, Sara Calhim, and Tim R. Birkhead, which appeared in issue 13, March 29, 2011, of *Proc Natl Acad Sci USA* (108:5325–5330; first published March 14, 2011; 10.1073/pnas.1009059108).

The authors note that, due to a printer’s error, the affiliation for Kate L. Durrant should instead appear as School of Biology, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom. The corrected author and affiliation lines appear below. The online version has been corrected.

**Simone Immler^{a,b,1,2}, Scott Pitnick^c, Geoff A. Parker^d,
Kate L. Durrant^e, Stefan Lüpold^c, Sara Calhim^f, and
Tim R. Birkhead^g**

^aDepartment of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield S10 2TN, United Kingdom; ^bDepartment of Evolutionary Biology, Evolutionary Biology Centre, Uppsala University, 75236 Uppsala, Sweden; ^cDepartment of Biology, Syracuse University, Syracuse, NY 13244; ^dDepartment of Evolution, Ecology and Behaviour, Institute of Integrative Biology, University of Liverpool, Liverpool L69 7ZB, United Kingdom; ^eSchool of Biology, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom; and ^fDepartment of Biology, Queen’s University, Kingston, ON, Canada K7L 3N6

www.pnas.org/cgi/doi/10.1073/pnas.1105524108

NEUROSCIENCE

Correction for “Conditional reduction of adult neurogenesis impairs bidirectional hippocampal synaptic plasticity,” by Federico Massa, Muriel Koelh, Theresa Wiesner, Noelle Grosjean, Jean-Michel Revest, Pier-Vincenzo Piazza, Djoher Nora Abrous, and Stéphane H. R. Oliet, which appeared in issue 16, April 19, 2011, of *Proc Natl Acad Sci USA* (108:6644–6649; first published April 4, 2011; 10.1073/pnas.1016928108).

The authors note that the author name Muriel Koelh should have appeared as Muriel Koehl. The corrected author line appears below. The online version has been corrected.

**Federico Massa, Muriel Koehl, Theresa Wiesner, Noelle
Grosjean, Jean-Michel Revest, Pier-Vincenzo Piazza,
Djoher Nora Abrous, and Stéphane H. R. Oliet**

www.pnas.org/cgi/doi/10.1073/pnas.1105938108